## (19) World Intellectual Property Organization

International Bureau





(43) International Publication Date 4 August 2005 (04.08.2005)

### **PCT**

# (10) International Publication Number WO 2005/070976 A1

(51) International Patent Classification<sup>7</sup>: C08F 210/16, 2/34

(21) International Application Number:

PCT/GB2005/000157

(22) International Filing Date: 14 January 2005 (14.01.2005)

(25) Filing Language:

English

(26) Publication Language:

English

(30) Priority Data:

0401348.8

22 January 2004 (22.01.2004) GF

- (71) Applicant (for all designated States except US): BP CHEMICALS LIMITED [GB/GB]; Chertsey Road, Sunbury on Thames, Middlesex TW16 7BP (GB).
- (72) Inventors; and
- (75) Inventors/Applicants (for US only): JACOBSEN, Grant, Berent [GB/BE]; Ringlaan 59, B-3080 Tervuren (BE). LALANNE-MAGNE, Claudine, Viviane [FR/FR]; 32 bis, Boulevard Pablo Neruda, F-13920 Saint Mitre les Remparts (FR). SERE PEYRIGAIN, Pierre [FR/FR]; 3, Aire des Cystes, F-13800 Istres (FR).
- (74) Agent: HAWKINS, David, George; O & D Trading Limited, Patents and Agreements, Compass Point, 79-87 Kingston Road, Staines, Middlesex TW18 1DT (GB).
- (81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD,

MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

#### Declarations under Rule 4.17:

- as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for the following designations AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NA, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SM, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, UZ, VC, VN, YU, ZA, ZM, ZW, ARIPO patent (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, MC, NL, PL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG)
- of inventorship (Rule 4.17(iv)) for US only

### Published:

- with international search report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: POLYMERISATION PROCESS

(57) Abstract: A process for the copolymerisation of ethylene and an  $\alpha$ -olefin having 7 to 10 carbon atoms in a fluidised bed gas phase reactor in the presence of a single site polymerisation catalyst comprises operating the process in condensed mode and wherein the amount of said  $\alpha$ -olefin is maintained below that at which substantial condensation in the reactor occurs. The preferred  $\alpha$ -olefin is 1-octene and the preferred single site polymerisation catalyst is a metallocene complex. By use of the process conditions of the present invention, higher  $\alpha$ -olefins may be successfully employed in a gas phase process provided the amount of higher  $\alpha$ -olefin comonomer is maintained below that at which substantial condensation occurs: Such operation is also dependent upon the operation temperature and the boiling point of the higher  $\alpha$ -olefin and the process is particularly advantageous when performed in the presence of catalysts able to incorporate high levels of comonomers at low comonomer inventories.





2